


Mercury in Tomales Bay

Data & Statistical Analysis for Training Academy

Carrie M. Austin, P.E.


San Francisco Bay Regional Water Board

**Tomales Bay
watershed**



**San
Francisco
Bay**

**Gambonini
Mercury Mine
drains to
Walker Creek**



Good reason to believe Tomales Bay is polluted by mercury, because of Gambonini mercury mine on Walker Creek.

Data set is focused on the Walker Creek delta, shown in red (red = mercury & elevated concentrations)

Several previous studies



Mercury accumulation and attenuation at a rapidly forming delta with a point source of mining waste

Bryce E. Johnson^{a,*}, Bradley K. Esser^b, Dyan C. Whyte^c, Priya M. Ganguli^c, Carrie M. Austin^c, James R. Hunt^a

^a Civil and Environmental Engineering, University of California, Berkeley, CA 94720, USA

^b Chemical Sciences Division, 7000 East Avenue, Lawrence Livermore National Laboratory, Livermore CA 94550, USA

^c California Regional Water Quality Control Board, 1515 Clay St., Suite 1400, Oakland, CA 94612, USA

ARTICLE INFO

Article history:

Received 14 December 2008

Received in revised form 6 May 2009

Accepted 12 May 2009

Available online xxxxx

Keywords:

Sediments

Mercury

Methylmercury

Contaminant accumulation

Bioaccumulation

Biota


Natural attenuation

Tomales Bay


ABSTRACT

The Walker Creek intertidal delta of Tomales Bay, California is impacted by a former mercury mine within the watershed. Eleven short sediment cores (10 cm length) collected from the delta found monomethylmercury (MMHg) concentrations ranging from 0.3 to 11.4 ng/g (dry wt.), with lower concentrations occurring at the vegetated marsh and upstream channel locations. Algal mats common to the delta's sediment surface had MMHg concentrations ranging from 7.5 to 31.5 ng/g, and the top 1 cm of sediment directly under the mats had two times greater MMHg concentrations compared to adjacent locations without algal covering. Spatial trends in resident biota reflect enhanced MMHg uptake at the delta compared to other bay locations. Eighteen sediment cores, 1 to 2 m deep, collected from the 1.2 km² delta provide an estimate of a total mercury (Hg) inventory of 2500 ± 500 kg. Sediment Hg concentrations ranged from pre-mining background conditions of approximately 0.1 µg/g to a post-mining maximum of 5 µg/g. Sediment accumulation rates were determined from three sediment cores using measured differences of ²¹⁰Pb activity. We estimate a pre-mining Hg accumulation of less than 20 kg/yr, and a period of maximum Hg accumulation in the 1970s and 1980s with loading rates greater than 50 kg/yr, corresponding to the failure of a tailings dam at the mine site. At the time of sampling (2003) over 40 kg/yr of Hg was still accumulating at the delta, indicating limited recovery. We attribute observed spatial evolution of elevated Hg levels to ongoing inputs and sediment reworking, and estimate the inventory of the anthropogenic fraction of total Hg to be at least 1500 ± 300 kg. We suggest ongoing sediment inputs and methylation at the deltaic surface support enhanced mercury levels for resident biota and transfer to higher trophic levels throughout the Bay.

© 2009 Elsevier B.V. All rights reserved.



Average surface sediment (upper 5 cm) mercury
Error bars are two times the standard deviation.
3 to 5 samples per location.



We focused our sediment sampling (brown) at opposite ends of Tomales Bay, shown in these insets, at Walker & Lagunitas deltas, expecting them to be most different (and most biologically important)

Sediment Sampling

PLAN

- $n = 8$ for each of Walker Creek delta and Lagunitas Creek delta at head of bay, additional samples from along the bay
- Each sample composited in field from 4 locations 1 meter apart
- Each delta sample collected in replicate within 100 meters (submitted individually to lab, then average results)


REALITY

- field error, $n = 6$, not 8, at Lagunitas

Total mercury in sediment (2009)

Walker Creek delta is obviously different, and
“other” is similar to Lagunitas Creek delta

Is the difference statistically significant?



ACTIVITY:

Compare total mercury in sediment from Walker Creek delta to “other” and “Lagunitas Creek delta” sites combined

- Import dataset (comma separators):
www.ics.uci.edu/~jutts/data/WalkerDelta.txt
- Recode variables, give new name, use “Lagunitas Delta” = “other”
- Do boxplots (by groups) of mercury
- Do independent sample t-test to compare mercury; look at CI for difference.
- Create new variable: $\log(\text{mercury})$ and repeat.